

Disseminated Intravascular Coagulation (DIC) Profile

Order Name: DIC PR Test Number: 1500855 Revision Date: 08/26/2014

TEST NAME		METHO	DOLOGY	LOINC CODE
Disseminated Intravascular Coagulation (DIC) Profile				
SPECIMEN REQUIREM	IENTS			
Specimen	Specimen Volume (min)	Specimen Type	Specimen Container	Transport Environment
Preferred	See Instructions	See Special Instructions	Sodium Citrate 3.2% (Blue Top) and EDTA (Lavender Top)	See Instructions
Instructions	Coagulation- Two: 2.7mL Sodium Citrate 3.2% (Blue Top) Tubes. (Ambient whole blood or frozen aliquots.) Please indicate anticoagulant therapy. Each 2.7mL Sodium Citrate 3.2% (Blue Top) tube must be filled to the proper level, no hemolysis. Improperly filled tubes can give erroneous results. Whole blood must be transported to lab immediately. If testing cannot be started within 4 hours of collection the specimen must be double spun then 1.5mL plasma aliquot from each tube into individual plastic aliquot tubes and freeze. Do not pool aliquots together! Specimen Stability: Un-Frozen specimens are only good for 4 hours. Hematology- One: 4mL (1mL) EDTA (Lavender Top) Whole Blood (Room Temperature)			
	Note: IPF level will not be reported	• • • •		
GENERAL INFORMATION				
Expected TAT	1 Day			

Expected TAT
1 Day

CPT Code(s)
85055, 85045, 85049, 85379, 85384, 85610, 85730

Service Provided By
Isophysical Composition of the com