

Tickborne Disease Panel

Instructions

Order Name: TICK PANEL Test Number: 5571750 Revision Date: 04/29/2024

TEST NAME		ME ⁻	THODOLOGY	LOINC CODE	
Tickborne Disease Panel					
SPECIMEN REQUIREMENTS					
Specimen	Specimen Volume (min)	Specimen Type	Specimen Container	Transport Environment	
Preferred	3 mL (1.5)	Serum	Clot Activator SST	Refrigerated or Frozen	

Allow specimen to clot completely at room temperature. Separate serum or plasma from cells ASAP or within 2 hours of collection.

Specimen Stability: Room Temperature: Unacceptable, Refrigerated: 2 Weeks, Frozen 30 Days

GENERAL INFORMATION			
Testing Schedule	Assay Dependant		
Expected TAT	3-5 Days		
Clinical Use	E. chaffeensis - A tick-borne disease common to the OK/ARK/MO/KS area. Peak titers will occur at 6 weeks after onset. Only 22% will be positive after 1 week, 68% after 2 weeks and 100% after 4 weeks. Rocky Mountain Spotted Fever - To assist in the diagnosis of RMSF. Patients may not seroconvert until 10 days after onset of illness. Tularemia Antibody - Diagnosis of exposure to Francisella tularensis.		
CPT Code(s)	86666x2, 86757x2, 86668x2		
Service Provided By	labcorp Oklahoma, Inc.		