


**Glucose Tolerance (5 Hour) w/o Urine**

Order Name: **GTT 5 HR**  
Test Number: 2002425  
Revision Date: 05/20/2021

TEST NAME	METHODOLOGY	LOINC CODE
Glucose Fasting	<a href="#">Hexokinase</a>	1558-6
Glucose 1 Hour Tolerance	<a href="#">Hexokinase</a>	20438-8
Glucose 2 Hour Total	<a href="#">Hexokinase</a>	20436-2
Glucose 3 Hour Tolerance	<a href="#">Hexokinase</a>	20437-0
Glucose 4 Hour Tolerance	<a href="#">Hexokinase</a>	26541-3
Glucose 5 Hour Tolerance	<a href="#">Hexokinase</a>	26543-9

SPECIMEN REQUIREMENTS				
Specimen	Specimen Volume (mL)	Specimen Type	Specimen Container	Transport Environment
Preferred	1 mL (0.5)	Plasma	Sodium fluoride (gray top)	Refrigerated
Alternate 1	1 mL (0.5)	Plasma	Lithium Heparin PST (Light Green Top)	Refrigerated
Alternate 2	1 mL (0.5)	Serum	Clot Activator SST	Refrigerated
Instructions	Fasting overnight and during test. Patient may have water. Draw fasting specimen. Adults: Give 75 grams of glucola. Children: Adjusted amount of glucola to be calculated by lab. Call (918) 744-2500. Stability: Ambient 8 hours, Refrigerated 7 days, Frozen 1 month Fluoride preserved plasma samples are stable at room temperature for 24 hours			

GENERAL INFORMATION	
Testing Schedule	Daily
Expected TAT	1-2 days
Clinical Use	There are no established guidelines, reference ranges or criteria, for the evaluation of a three hour, four hour or five hour specimen following a five hour glucose challenge.
Notes	For more information on this test, access our "Specialized Tests" section.
CPT Code(s)	82951, 82952x3
Service Provided By	 <b>labcorp</b> Oklahoma, Inc.