

Zinc, Serum or Plasma

Order Name: ZINC
Test Number: 3603800
Revision Date: 12/12/2022

TEST NAME	METHODOLOGY	LOINC CODE
Zinc, Serum or Plasma	Inductively-Coupled Plasma/Mass Spectrometry	
SPECIMEN REQUIREMENTS		

SPECIMEN REQUIREMENTS						
		On a simon Toma	On a signature On a taige an	Transport For incorporat		
Specimen	Specimen Volume (min)	Specimen Type	Specimen Container	Transport Environment		
Preferred	2 mL (0.6)	Serum	Royal Blue - No Additive Clot	Ambient / Refrigerated		
			(Trace Element Free)			
Alternate 1	2 mL (0.6)	Plasma	Royal Blue - EDTA (Trace Element	Ambient / Refrigerated		
			Free)			
Instructions	Notes: 0.6 mL (Note: This vol	Notes: 0.6 mL (Note: This volume Does NOT allow for repeat testing.)				
Specimen Type: Royal blue-top (EDTA) tube OR royal blue-top without EDTA						
	Specimen Storage: Maintain specimen at room temperature.					
	Specimen Collection: Separate serum from cells within 45 minutes of collection and transfer to a certified metal-free transport tube (PeopleSoft No.					
	111166) for shipment to the laboratory. Plasma may be separated immediately and transferred to a certified metal-free plastic transport tube					
	(PeopleSoft No. 111166 for shipment to the laboratory.					
	Specimen Stability: Ambient: 14 days, Refrigerated : 14 days, Frozen: 14 days					

GENERAL INFORMATION	
Expected TAT	2 - 4 days within one working day.
Clinical Use	Serum or plasma measurements are best for assessment of deficiency over short periods of time. For acute toxicity, zinc in urine may be a more reliable indicator of exposure.
Notes	Labcorp Test Code: 001800
CPT Code(s)	84630
Service Provided By	labcorp Oklahoma, Inc.