

Order Name: LIDOCAINE Test Number: 3603900 Revision Date: 12/12/2022

TEST NAME			METHODOLOGY			
Lidocaine (Xylocaine)		Immunoassay (IA)		3714-3		
SPECIMEN REQUIREMENTS						
Specimen	Specimen Volume (min)	Specimen Type	Specimen Container	r	Transport Environment	
Preferred	1 mL (0.3)	Serum	Clot Activator (Red	l Top, No-Gel)	Room Temperature	
Alternate 1	1 mL (0.3)	Plasma	Sodium Heparin (G Gel)	reen Top / No-	Room Temperature	
Instructions	<ul> <li>Notes: 0.3 mL (Note: This volume Does NOT allow for repeat testing.)</li> <li>Specimen Type: Red-top tube OR green-top (heparin) tube. DO NOT USE A GEL-BARRIER TUBE. The use of a gel-barrier tubes is not recommended due to slow absorption of the drug by the gel. Depending on the specimen volume and storage time, the decrease in drug level due to absorption may be clinically significant.</li> <li>Container Detail: No Gel</li> <li>Specimen Storage: Room Temperature</li> <li>Specimen Collection: Transfer separated serum or plasma to a plastic transport tube. Draw specimens 12 hours after initiating therapy for arrhythmia prophylaxis, then every 24 hours thereafter. Obtain specimens every 12 hours when cardiac or hepatic insufficiency exists.</li> <li>Specimen Stability: Ambient: 14 days, Refrigerated : 14 days, Frozen: 14 days</li> </ul>					
GENERAL INFORMAT	TION					

GENERAL INFORMATION		
Expected TAT	2 - 3 days Saturday thru Sunday on 1st shift; reported within one working day.	
Notes	Labcorp Test Code: 007013	
CPT Code(s)	80176	
Service Provided By	Oklahoma, Inc.	