

Flecainide (Tambocor)

Order Name: **TAMBOCOR**Test Number: 3618200
Revision Date: 12/12/2022

TEST NAME		METHODOLOGY LOINC COD		LOINC CODE
Flecainide (Tambocor)		<u>Ligui</u>	Chromatography/Tandem Mass Spectromet	<u>ry</u> 3638-4
SPECIMEN REQUIREMENTS				
Specimen	Specimen Volume (min)	Specimen Type	Specimen Container	Transport Environment
Preferred	1 mL (0.2 mL)	Serum	Clot Activator (Red Top, No-Gel)	Refrigerated
Alternate 1	1 mL (0.2 mL)	Plasma	Lithium Heparin (Dark Green Top / No-Gel)	Refrigerated
Instructions	Specimen Type: Red-top (no additive) tube or green-top (heparin) tube; gel-barrier tubes not recommended Specimen Storage: Refrigerated Specimen Collection: Transfer separated serum or plasma to a plastic transport tube. Draw immediately prior to next dose unless otherwise instructed. Specimen Stability: Ambient: 3 days, Refrigerated: 14 days, Frozen: 14 days			

GENERAL INFORMATION		
Expected TAT	2 - 4 days working day.	
Clinical Use	Monitoring the flecainide concentration is used to assure compliance and avoid toxicity of this cardiac drug used to treat ventricular tachcardia and premature cardiac contractions.	
Notes	Labcorp Test Code: 085662	
CPT Code(s)	80181	
Service Provided By	labcorp Oklahoma, Inc.	