

Flecainide (Tambocor)

Order Name: **TAMBOCOR**Test Number: 3618200
Revision Date: 12/12/2022

| TEST NAME | | METHO | DDOLOGY | LOINC CODE | |
|-----------------------|---|---------------|---|-----------------------|--|
| Flecainide (Tambocor) | cor) Liqui | | Chromatography/Tandem Mass Spectrometi | <u>ry</u> 3638-4 | |
| SPECIMEN REQUIREMENTS | | | | | |
| Specimen | Specimen Volume (min) | Specimen Type | Specimen Container | Transport Environment | |
| Preferred | 1 mL (0.2 mL) | Serum | Clot Activator (Red Top, No-Gel) | Refrigerated | |
| Alternate 1 | 1 mL (0.2 mL) | Plasma | Lithium Heparin (Dark Green Top / No-Gel) | Refrigerated | |
| Instructions | Specimen Type: Red-top (no additive) tube or green-top (heparin) tube; gel-barrier tubes not recommended Specimen Storage: Refrigerated Specimen Collection: Transfer separated serum or plasma to a plastic transport tube. Draw immediately prior to next dose unless otherwise instructed. Specimen Stability: Ambient: 3 days, Refrigerated: 14 days, Frozen: 14 days | | | | |

| GENERAL INFORMATION | | |
|---------------------|---|--|
| Expected TAT | 2 - 4 days working day. | |
| Clinical Use | Monitoring the flecainide concentration is used to assure compliance and avoid toxicity of this cardiac drug used to treat ventricular tachcardia and premature cardiac contractions. | |
| Notes | Labcorp Test Code: 085662 | |
| CPT Code(s) | 80181 | |
| Service Provided By | labcorp Oklahoma, Inc. | |