

Acetylcholine Receptor Blocking Antibody

Order Name: ACETY BLK
Test Number: 5500020
Revision Date: 01/02/2024

TEST NAME			METHODOLOGY	LOINC CODE	
Acetylcholine Receptor Blocking Antibody			See Test Notes	11561-8	
SPECIMEN REQUIREMENTS					
Specimen	Specimen Volume (min)	Specimen Type	Specimen Container	Transport Environment	
Preferred	1 mL (0.3)	Serum	Clot Activator SST	Refrigerated	
Alternate 1	1 mL (0.3)	Serum	Clot Activator (Red Top, No-Gel)	Refrigerated	
Instructions	Notes: 0.3 mL (Note: This volume Does NOT allow for repeat testing.) Specimen Type: Red-top tube or gel-barrier tube Specimen Storage: Refrigerate Specimen Collection: If red-top tube is used, transfer separated serum to a plastic transport tube. Specimen Stability: Ambient: 13 days, Refrigerated: 13 days, Frozen: 13 days				

GENERAL INFORMATION	
Expected TAT	3 - 7 days
Clinical Use	Blocking antibodies are detected in approximately 50% of generalized myasthenia gravis patients and are detectable in the absence of binding antibodies in approximately 1% of myasthenia gravis patients.
Notes	This assay measures antibodies that inhibit the binding of radiolabeled alpha-bungarotoxin (aBTX) to solublized muscle AChR.
CPT Code(s)	86042
Service Provided By	labcorp Oklahoma, Inc.