

Glutamic Acid Decarboxylase-65 Autoantibodies

Order Name: ANTI GAD
Test Number: 5592950
Revision Date: 02/27/2023

TEST NAME		METHODOLOGY		LOINC CODE
Glutamic Acid Decarboxylase-65 Autoantibodies		Enzyme	e-Linked Immunosorbent Assay (ELISA)	56540-8
SPECIMEN REQUIREM	MENTS			
Specimen	Specimen Volume (min)	Specimen Type	Specimen Container	Transport Environment
Preferred	1 mL (0.4 mL)	Serum	Clot Activator SST	Room Temperature
Alternate 1	1 mL (0.4 mL)	Serum	Clot Activator (Red Top, No-Gel)	Room Temperature
Instructions	Notes: 0.4 mL (Note: This volume Does NOT allow for repeat testing.) Specimen Type: Red-top tube or gel-barrier tube Specimen Storage: Room temperature Specimen Collection: If a red-top tube is used, transfer separated serum to a plastic transport tube. Avoid hemolysis. Special Instructions: This test may exhibit interference when sample is collected from a person who is consuming a supplement with a high dose of biotin (also termed as vitamin B7 or B8, vitamin H, or coenzyme R). It is recommended to ask all patients who may be indicated for this test about biotin supplementation. Patients should be cautioned to stop biotin consumption at least 72 hours prior to the collection of a sample Specimen Stability: Ambient: 14 days, Refrigerated: 14 days, Frozen: 14 days			

GENERAL INFORMATION		
Expected TAT	3 - 4 days	
Notes	Labcorp Test Code: 143008	
CPT Code(s)	86341	
Service Provided By	labcorp Oklahoma, Inc.	