

## Striated Muscle Antibody Screen with Titer

Order Name: **STRIAT AB**  
 Test Number: 3805400  
 Revision Date: 03/10/2026

TEST NAME	METHODOLOGY	LOINC CODE
Striated Muscle Antibody Screen	<u>Semi-Quantitative Indirect Fluorescent Antibody</u>	5372-8

### SPECIMEN REQUIREMENTS

Specimen	Specimen Volume (min)	Specimen Type	Specimen Container	Transport Environment
Preferred	1 mL (0.5 mL)	Serum	Clot Activator SST	Refrigerated
Alternate 1	1 mL (0.5 mL)	Serum	Clot Activator (Red Top, No-Gel)	Refrigerated

**Instructions**

**Minimum Volume:** 0.5 mL (Note: This volume does not allow for repeat testing.)  
**Specimen Type:** Gel-barrier tube or Red-top tube  
**Specimen Storage:** Refrigerated  
**Specimen Collection:** Separate serum from cells ASAP or within 2 hours of collection  
**Specimen Stability:** Ambient: 14 days, Refrigerated : 14 days, Frozen: 14 days (Freeze/thaw cycles Stable x3)  
**Causes for Rejection:** Hemolyzed or lipemic serum

**If being collected for Lambert-Eaton Syndrome Panel, Please collect Serum from Red No-Gel Clot Tube.**

### GENERAL INFORMATION

Expected TAT	2 - 4 days
Clinical Use	This test is recommended for differential evaluation of neuromuscular junction diseases including myasthenia gravis (MG). This test will report a Negative Screen and if Positive will report out the Titer. <b>Reference Interval</b> Negative: <1:100
CPT Code(s)	86255

Service Provided By  **labcorp**  
Oklahoma, Inc.