

**Striated Muscle Antibody Screen with Titer**

 Order Name: **STRIAT AB**

Test Number: 3805400

Revision Date: 02/10/2023

TEST NAME	METHODOLOGY	LOINC CODE
Striated Muscle Antibody Screen	Semi-Quantitative Indirect Fluorescent Antibody	5372-8

**SPECIMEN REQUIREMENTS**

Specimen	Specimen Volume (min)	Specimen Type	Specimen Container	Transport Environment
Preferred	<b>1 mL (0.5 mL)</b>	<b>Serum</b>	<b>Clot Activator SST</b>	<b>Refrigerated</b>
Alternate 1	<b>1 mL (0.5 mL)</b>	<b>Serum</b>	<b>Clot Activator (Red Top, No-Gel)</b>	<b>Refrigerated</b>
Instructions	<b>Minimum Volume:</b> 0.5 mL (Note: This volume does not allow for repeat testing. <b>Specimen Type:</b> Gel-barrier tube or Red-top tube <b>Specimen Storage:</b> Refrigerated <b>Specimen Collection:</b> Separate serum from cells ASAP or within 2 hours of collection <b>Specimen Stability:</b> Ambient: 14 days, Refrigerated : 14 days, Frozen: 14 days (Freeze/thaw cycles Stable x3) <b>Causes for Rejection:</b> Hemolyzed or lipemic serum			
<b>If being collected for Lambert-Eaton Syndrome Panel, Please collect Serum from Red No-Gel Clot Tube.</b>				

**GENERAL INFORMATION**

Expected TAT	2 - 4 days
Clinical Use	This test is recommended for differential evaluation of neuromuscular junction diseases including myasthenia gravis (MG). This test will report a Negative Screen and if Positive will report out the Titer. <b>Reference Interval</b> Negative: <1:100
Notes	Labcorp Test Code: 160184
CPT Code(s)	86255
Service Provided By	 <b>labcorp</b> Oklahoma, Inc.