METHODOL OCY



82610

labcorp
Oklahoma, Inc.

Cystatin C

CPT Code(s)

Service Provided By

Order Name: CYSTATIN C
Test Number: 3623775
Revision Date: 10/29/2025

TEST NAME		MET	HODOLOGY	LOINC CODE	
Cystatin C		Turk	Turbidometric		
SPECIMEN REQUIRE	EMENTS				
Specimen	Specimen Volume (min)	Specimen Type	Specimen Container	Transport Environment	
Preferred	1 mL (0.5 mL)	Serum	Clot Activator SST	Refrigerated	
Alternate 1	1 mL (0.5 mL)	Plasma	Sodium Heparin (Green Top / No- Gel)	- Refrigerated	
Instructions	Collect blood in SST Clot tube of Centrifuge aliquot Serum or Plaspecimen Stability: Room temp	Dietary Instructions: Overnight fasting is preferred. Collect blood in SST Clot tube or Sodium Heparin Green-top tube. Allow Serum to clot for 15 minutes. Centrifuge aliquot Serum or Plasma for 10min and separate wihtin 2 hours and keep refrigerated. Specimen Stability: Room temperature: 24 Hours, Refrigerated: 7 Days, Frozen: 90 Days.			
GENERAL INFORMA					
Testing Schedule	Mon-Fri				
Expected TAT	3-4 Days				
Clinical Use	, , ,	Cystatin C is a highly sensitive and specific marker of glomerular filtration rate (renal function). Cystatin C is independent of muscle mass, age, and body mass index. Cystatin C is also used to assess renal allograft function.			
Notes	This test will order and result a Cystatin C Estimated GFR as shown below.				
	ORDER CODE	REFLEX NAME	RESULT CODE	RESULT LONG NAME	