

Cystatin C

Order Name: **CYSTATIN C**
 Test Number: 3623775
 Revision Date: 10/29/2025

TEST NAME	METHODOLOGY	LOINC CODE
Cystatin C	Turbidometric	33863-2

SPECIMEN REQUIREMENTS

Specimen	Specimen Volume (min)	Specimen Type	Specimen Container	Transport Environment
Preferred	1 mL (0.5 mL)	Serum	Clot Activator SST	Refrigerated
Alternate 1	1 mL (0.5 mL)	Plasma	Sodium Heparin (Green Top / No-Gel)	Refrigerated

Instructions

Dietary Instructions: **Overnight fasting is preferred.**
 Collect blood in SST Clot tube or Sodium Heparin Green-top tube. Allow Serum to clot for 15 minutes.
 Centrifuge aliquot Serum or Plasma for 10min and separate within 2 hours and keep refrigerated.
 Specimen Stability: Room temperature: 24 Hours, Refrigerated: 7 Days, Frozen: 90 Days.

GENERAL INFORMATION

Testing Schedule	Mon-Fri
Expected TAT	3-4 Days
Clinical Use	Cystatin C is a highly sensitive and specific marker of glomerular filtration rate (renal function). Cystatin C is independent of muscle mass, age, and body mass index. Cystatin C is also used to assess renal allograft function.
Notes	This test will order and result a Cystatin C Estimated GFR as shown below.

ORDER CODE	REFLEX NAME	RESULT CODE	RESULT LONG NAME
3623778	.Cystatin C Estimated GFR	3623888	Cystatin C eGFR

CPT Code(s)	82610
Service Provided By	 labcorp Oklahoma, Inc.