St. John Health System Lab Catalog

Cryptococcus Antigen Screen - Serum

Order Name: CRYPTO AG
Test Number: 6002175
Revision Date: 02/21/2020

TEST NAME		METHO	DDOLOGY	LOINC CODE	
Cryptococcus Antigen Screen - Serum		Enzym	e Immunoassay	31790-9	
SPECIMEN REQUIREMENTS					
Specimen	Specimen Volume (min)	Specimen Type	Specimen Container	Transport Environment	
Preferred	4 mL (1)	Serum	Clot Activator (Red Top, No-Gel)	Refrigerated	
Alternate 1	4 mL (1)	Serum	Clot Activator SST (Gold Top)	Refrigerated	
Instructions	Cleanse venipuncture site and avoid hemolyzed specimens. Allow specimen to clot and aliquot 4mL(1mL) of serum to a standard transport tube. Store serum at 2-8 ? for up to 72 hours. Serum may be poured off and frozen at <-20 ? for up to 7 days prior to testing. Specimens should remain at 2-8 or -20 during transport. Unacceptable specimens: Hemolyzed specimens, specimens contaminated with gel, specimens frozen then thawed and refrozen, specimens stored at the incorrect temperature, or specimens contaminated.				

GENERAL INFORMATION		
Testing Schedule	Mon - Fri	
Expected TAT	1 Day	
Clinical Use	Detects presence of Cryptococcus antigen in peripheral blood	
CPT Code(s)	87899 If the screen is positive, then the following test will be performed:	
	Order Code [6002300] Cryptococcus Antigen Titer	
	Result Code [6002300] Cryptococcus Ag Titer	
	Cpt Code 86406	
Service Provided By	labcorp Oklahoma, Inc.	