


**Clam IgG**

Order Name: **CLAM IGG**  
Test Number: 5500507  
Revision Date: 12/14/2022

TEST NAME	METHODOLOGY	LOINC CODE
Clam IgG	Enzyme immunoassay (EIA)	

SPECIMEN REQUIREMENTS				
Specimen	Specimen Volume (min)	Specimen Type	Specimen Container	Transport Environment
Preferred	0.3 mL (0.1 mL)	Serum	Clot Activator SST	Room Temperature
Instructions	<p><b>Specimen Type:</b> Red-top tube or gel-barrier tube</p> <p><b>Specimen Storage:</b> Room temperature</p> <p><b>Specimen Collection:</b> Not Available</p> <p><b>Special Instructions:</b> Values obtained with different assay methods should not be used interchangeably in serial testing. It is recommended that only one assay method be used consistently to monitor each patient's course of therapy.</p> <p><b>Specimen Stability:</b> Ambient: Not Available, Refrigerated : Not Available, Frozen: Not Available</p>			

GENERAL INFORMATION	
Expected TAT	2 - 4 days
Clinical Use	The clinical utility of food-specific IgG tests has not been established. These tests can be used in special clinical situations to select foods for evaluation by diet elimination and challenge in patients who have food-related complaints. It should be recognized that the presence of food-specific IgG alone cannot be taken as evidence of food allergy and only indicates immunologic sensitization by the food allergen in question. This test should only be ordered by physicians who recognize the limitations of the test. The reference range listed on the report is the lower limit of quantitation for the assay.
Notes	Labcorp Test Code: 605119
CPT Code(s)	86001
Service Provided By	 <b>labcorp</b> Oklahoma, Inc.