

Diazepam and NordiazepamOrder Name: **DIAZEPAM**

Test Number: 3611375

Revision Date: 10/18/2017

TEST NAME	METHODOLOGY	LOINC CODE
Diazepam and Nordiazepam	Liquid Chromatography/Tandem Mass Spectrometry	

SPECIMEN REQUIREMENTS				
Specimen	Specimen Volume (mL)	Specimen Type	Specimen Container	Transport Environment
Preferred	1 mL (0.4 mL)	Serum	Clot Activator (Red Top, No-Gel)	Refrigerated
Instructions	<p>Specimen Type: Red-top tube, lavender-top (EDTA) tube OR green-top (heparin) tube. DO NOT USE A GEL-BARRIER TUBE. The use of gel-barrier tubes is not recommended due to slow absorption of the drug by the gel. Depending on the specimen volume and storage time, the decrease in drug level due to absorption may be clinically significant.</p> <p>Specimen Storage: Refrigerate</p> <p>Specimen Collection: Transfer separated serum or plasma to a plastic transport tube. Oral: peak; one hour after dose; I.V.: peak: 15 minutes after dose.</p> <p>Specimen Stability: Ambient: 1 day, Refrigerated : 14 days, Frozen: 14 days</p>			

GENERAL INFORMATION	
Expected TAT	2-4 Days
Notes	Labcorp Test Code: 007989
CPT Code(s)	80299
Service Provided By	 labcorp Oklahoma, Inc.