


Heparin Cofactor II

Order Name: **Heparin Cof II**
Test Number: 6905029
Revision Date: 12/12/2022

TEST NAME	METHODOLOGY	LOINC CODE
Heparin Cofactor II	Chromogenic	33987-9

SPECIMEN REQUIREMENTS				
Specimen	Specimen Volume (min)	Specimen Type	Specimen Container	Transport Environment
Preferred	2 mL (1)	Platelet Poor Plasma	Sodium Citrate 3.2% (Blue Top)	Frozen
Instructions	<p>Specimen Type: Blue-top (sodium citrate) tube</p> <p>Specimen Storage: Frozen</p> <p>Specimen Collection: CITRATED PLASMA SAMPLES SHOULD BE COLLECTED BY DOUBLE CENTRIFUGATION. Blood should be collected in a blue-top tube containing 3.2% buffered sodium citrate.[1] Evacuated collection tubes must be filled to completion to ensure a proper blood to anticoagulant ratio.[2,3] The sample should be mixed immediately by gentle inversion at least six times to ensure adequate mixing of the anti-coagulant with the blood. A discard tube is not required prior to collection of coagulation samples except when using a winged blood collection device (i.e. 'butterfly'), in which case a discard tube should be used.[4,5] When noncitrate tubes are collected for other tests, collect sterile and nonadditive (red-top) tubes prior to citrate (blue-top) tubes. Any tube containing an alternate anticoagulant should be collected after the blue- top tube. Gel-barrier tubes and serum tubes with clot initiators should also be collected after the citrate tubes. Centrifuge for 10 minutes and carefully remove 2/3 of the plasma using a plastic transfer pipette, being careful not to disturb the cells. Deliver to a plastic transport tube, cap, and recentrifuge for 10 minutes. Use a second plastic pipette to remove plasma, staying clear of the platelets at the bottom of the tube. Transfer the plasma into a LabCorp PP transpak frozen purple tube with screw cap (LabCorp 49482). FREEZE immediately and maintain frozen until tested. To avoid delays in turnaround time when requesting multiple tests on frozen samples, please submit separate frozen specimens for each test requested. Please print and use the Volume Guide for Coagulation Testing to ensure proper draw volume.</p> <p>Special Instructions: Testing performed at Esoterix Coagulation UY#300133</p> <p>Specimen Stability: Refrigerate up to 48 hours</p>			

GENERAL INFORMATION	
Expected TAT	5 - 8 days
Notes	Patient plasma is incubated with excess thrombin in the presence of dermatan sulfate. Residual thrombin activity reacts with chromogenic substrate and the color intensity is inversely proportional to heparin cofactor II levels. Heparin levels up to 1 U/mL ?Labcorp Test Code: 500187
CPT Code(s)	85130
Service Provided By	 labcorp Oklahoma, Inc.