St. John Health System Lab Catalog

CT/GC/TV Amplified-Cervix (send out)

Order Name: CT/GC/TV Cervix

Test Number: 5195343
Revision Date: 06/25/2024

TEST NAME		MET	HODOLOGY	LOINC CODE
CT/GC/TV Amplified-Cervix (send out)		Nuc	eic acid amplification (NAA)	See Notes
SPECIMEN REQUIREM	MENTS			
Specimen	Specimen Volume (min)	Specimen Type	Specimen Container	Transport Environment
Preferred	1	Endocervical swab	Cobas(R) PCR Female Swab Sample Kit	Room Temperature
Instructions	Aptima (Gen-probe) Endocx/ureth swab - cobas(R) PCR Female Swab Sample Kit Collection: Collecting an endocervical swab specimen for use in NAAT testing procedures involves the use of a speculum and includes the use of two swabs one for initial cleaning of the cervix to remove excess mucus from the cervical os and surround mucosa (this swab should be discarded), and a second for specimen collection. 1) Use one of the swabs in the collection kit to clean the mucous from the cervix. 2) Discard the cleaning swab. 3) Use a second swab to collect the specimen by insertion of the swab into the endocervical canal followed by gentle rotation of the swab. 4) Withdraw the collection swab while avoiding contact with the vaginal mucosa. 5) Place the swab into the PCR media tube and break the shaft at the score line. 6) Do not submit both swabs. 7) Tightly and securely seal the PCR media tube. 8) Mixed to ensure the collected specimen has been thoroughly exposed to the transport media which contains nucleic acid stabilizing agents. Storage Instructions: Room temperature Stability: Room temperature: 365 days Refrigerated: 365 days			

GENERAL INFORMATION		
Expected TAT	Assay Dependent	
Notes	Amplified TV-Cervix 70166-4	
	Amplified GC-Cervix 50388-8	
	Amplified CT-Cervix 21190-4	
CPT Code(s)	87491; 87591; 87661	
Service Provided By	Oklahoma, Inc.	

Cause for Rejection: Contamination, Bloody; 2 swabs in tube; less than 2 mL liquid; no swab inside the tube or inverted swabs