

Factor 5 (V) Assay

 Order Name: **FACTOR 5**

Test Number: 1501050

Revision Date: 10/08/2024

TEST NAME	METHODOLOGY	LOINC CODE
Factor 5 (V) Assay	Clot Detection	3193-0

SPECIMEN REQUIREMENTS

Specimen	Specimen Volume (min)	Specimen Type	Specimen Container	Transport Environment				
Preferred	See Instructions	Plasma	Sodium Citrate 3.2% (Blue Top)	Frozen				
Instructions	Please indicate anticoagulant therapy. Collect Two 2.7mL Sodium Citrate 3.2% (Blue Top) tubes <p>Each 2.7mL Sodium Citrate 3.2% (Blue Top) tube must be filled to the proper level, no hemolysis. Improperly filled tubes can give erroneous results. Whole blood must be transported to lab immediately. If testing cannot be started within 4 hours of collection the specimen must be double spun then 1.5mL plasma aliquot from each tube into individual plastic aliquot tubes and freeze. Do not pool aliquots together!</p>							

GENERAL INFORMATION

Testing Schedule	Tue and/or Thur - Day Shift
Expected TAT	1-7 Days
Stat TAT	M-F both shifts stat only
Clinical Use	This assay measures the clotting ability of Factor 5. This assay is used to aid in the diagnosis of coagulation factor deficiencies that may present with epistaxis, easy bruising, or menorrhagia
CPT Code(s)	85220
Service Provided By	 labcorp Oklahoma, Inc.