

**Disseminated Intravascular Coagulation (DIC) Profile**

Order Name: **DIC PR**  
 Test Number: 1500855  
 Revision Date: 08/26/2014

TEST NAME	METHODOLOGY	LOINC CODE
Disseminated Intravascular Coagulation (DIC) Profile		

SPECIMEN REQUIREMENTS				
Specimen	Specimen Volume (min)	Specimen Type	Specimen Container	Transport Environment
Preferred	<a href="#">See Instructions</a>	<a href="#">See Special Instructions</a>	<b>Sodium Citrate 3.2% (Blue Top) and EDTA (Lavender Top)</b>	<a href="#">See Instructions</a>
Instructions	<b>Coagulation-</b> Two: 2.7mL Sodium Citrate 3.2% (Blue Top) Tubes. (Ambient whole blood or frozen aliquots.)  <b>Please indicate anticoagulant therapy.</b> Each 2.7mL Sodium Citrate 3.2% (Blue Top) tube must be filled to the proper level, no hemolysis. Improperly filled tubes can give erroneous results. <b>Whole blood must be transported to lab immediately.</b> <b>If testing cannot be started within 4 hours of collection the specimen must be double spun then 1.5mL plasma aliquot from each tube into individual plastic aliquot tubes and freeze.</b> <b>Do not pool aliquots together!</b> Specimen Stability: Un-Frozen specimens are only good for 4 hours.			
<b>Hematology-</b> One: 4mL (1mL) EDTA (Lavender Top) Whole Blood (Room Temperature) Note: IPF level will not be reported on specimens > 24hrs old.				

GENERAL INFORMATION	
Expected TAT	1 Day
CPT Code(s)	85055, 85045, 85049, 85379, 85384, 85610, 85730
Service Provided By	 <b>labcorp</b> Oklahoma, Inc.