

**Vancomycin Peak**

Order Name: **VANCO PEAK**  
 Test Number: 4005900  
 Revision Date: 06/04/2021

| TEST NAME       | METHODOLOGY                             | LOINC CODE |
|-----------------|---|------------|
| Vancomycin Peak | Enzyme Multiplied Immunoassay Technique | 4090-7     |

**SPECIMEN REQUIREMENTS**

|              |   |               |  |                               |
|--------------|---|---------------|--|-------------------------------|
| Specimen     | Specimen Volume (min)   | Specimen Type | Specimen Container                           | Transport Environment         |
| Preferred    | <b>1 mL (0.5)</b>   | <b>Plasma</b> | <b>Lithium Heparin PST (Light Green Top)</b> | <b>Refrigerated or Frozen</b> |
| Alternate 1  | <b>1 mL (0.5)</b>   | <b>Serum</b>  | <b>Clot Activator SST</b>                    | <b>Refrigerated or Frozen</b> |
| Instructions | Peak: draw specimen 60 minutes after a 1 hour infusion.<br>Separate plasma/serum from cells/gel immediately.<br>Stability: Refrigerated 7 days, Frozen 14 days. |               |  |                               |

**GENERAL INFORMATION**

|                     |  |
|---------------------|--|
| Testing Schedule    | Daily  |
| Expected TAT        | 1-2 days   |
| Clinical Use        | Useful for optimizing drug dosage and assessing toxicity.  |
| CPT Code(s)         | 80202  |
| Service Provided By |  <b>labcorp</b><br>Oklahoma, Inc. |