

Tissue Transglutaminase IgG (IgG anti-tTG)

Order Name: **TISTRN IGG**
 Test Number: 5536025
 Revision Date: 12/12/2022

TEST NAME	METHODOLOGY	LOINC CODE
Tissue Transglutaminase IgG (IgG anti-tTG)	Enzyme immunoassay (EIA)	56537-4

SPECIMEN REQUIREMENTS

Specimen	Specimen Volume (min)	Specimen Type	Specimen Container	Transport Environment
Preferred	1 mL (0.5)	Serum	Clot Activator SST	Refrigerated
Instructions	Notes: 0.5 mL (Note: This volume Does NOT allow for repeat testing). Specimen Type: Red-top tube or gel-barrier tube Specimen Storage: Room temperature Specimen Collection: Specimen should be free of bacterial contamination, hemolysis, and lipemia. To avoid delays in turnaround time when requesting multiple tests on frozen samples, PLEASE SUBMIT SEPARATE FROZEN SPECIMENS FOR EACH TEST REQUESTED. Specimen Stability: Ambient: 14 days, Refrigerated : 14 days, Frozen: 14 days			

GENERAL INFORMATION

Expected TAT	2-3 Days
Clinical Use	Tissue Transglutaminase Antibody, IgG is useful in diagnosing gluten-sensitive enteropathies, such as celiac sprue, and an associated skin condition, dermatitis herpetiformis in patients who are IgA-deficient. The test also provides support for gluten-sensitive enteropathies beyond IgA the test.
Notes	Labcorp Test Code: 164988
CPT Code(s)	86364
Service Provided By	 labcorp Oklahoma, Inc.