

Anaplasma Phagocytophilum (HGA), IgG/IgM

Order Name: **A Phagocytoph Abs**
 Test Number: 5613565
 Revision Date: 08/29/2023

TEST NAME	METHODOLOGY	LOINC CODE
A Phagocytophilum Antibody, IgG	<u>Immunofluorescence Assay (IFA)</u>	23877-4
A Phagocytophilum Antibody, IgM	<u>Immunofluorescence Assay (IFA)</u>	23878-2

SPECIMEN REQUIREMENTS

Specimen	Specimen Volume (min)	Specimen Type	Specimen Container	Transport Environment
Preferred	0.5 mL (0.2 mL)	Serum	Clot Activator SST	Refrigerated
Instructions	Specimen Type: Red-top tube or gel-barrier tube Collection: If tube other than a gel-barrier is used, transfer separated serum to a labeled plastic transport tube. Do not freeze gel-barrier tube, pour off serum first. Stability: Room temperature: 14 days, Refrigerated: 14 days, Frozen: 14 days (Freeze/Thaw Cycle: 4x) ?Causes for Rejection: Hemolysis; lipemia; gross bacterial contamination; frozen gel-barrier tube			

GENERAL INFORMATION

Expected TAT	2- 4 days
Clinical Use	Acceptable test for acute or convalescent phase of infection from Anaplasma phagocytophilum . May be useful when PCR testing is not an option (eg, outside the 2 week window for acute phase).
CPT Code(s)	86666x2
Service Provided By	 labcorp Oklahoma, Inc.