

Mycoplasma genitalium, NAA, Urine

Order Name: **M genital, UR**
 Test Number: 5195311
 Revision Date: 04/21/2026

TEST NAME	METHODOLOGY	LOINC CODE
Mycoplasma genitalium, NAA, Urine	<u>Nucleic acid amplification (NAA)</u>	88226-6

SPECIMEN REQUIREMENTS

Specimen	Specimen Volume (min)	Specimen Type	Specimen Container	Transport Environment
Preferred	30 mL (2 mL)	Urine (See Inst.)	APTIMA Urine Transport Tube	Room Temperature

Instructions

Specimen: First-void urine in Aptima® urine transport
Volume: 2 mL of a 20 to 30 mL urine collection
Container: Aptima® urine transport
Collection: Urine specimen: The patient should not have urinated for at least one hour prior to specimen collection. Direct patient to provide a first-catch urine (approximately 20 to 30 mL of the initial urine stream) into a urine collection cup free of any preservatives. Collection of larger volumes of urine may result in specimen dilution that may reduce test sensitivity; lesser volumes may not adequately rinse organisms into the specimen. Female patients should not cleanse the labial area prior to providing the specimen. Add urine to the Aptima® Combo 2 urine collection device. The final volume must be between the two black lines on the device (about 2 mL).
Storage Instructions: Maintain specimen at room temperature or refrigerate (2°C to 30°C).
Causes for Rejection: **Not for Male Collection.**
Stability Room temperature 30 days (stability provided by manufacturer or literature reference)

GENERAL INFORMATION

Expected TAT	3-4 days
--------------	----------

Notes	RESULT CODE	RESULT NAME	LOINC
	6907667	M genitalia Ur	88226-6

CPT Code(s)	87563
-------------	-------

Service Provided By 